

AD-A138 683

PROGRESS AND PROBLEMS IN STORAGE RING FREE ELECTRON
LASERS(U) STANFORD UNIV CA HIGH ENERGY PHYSICS LAB

1/1

C BAZIN ET AL. FEB 83 HEPL-923 AFOSR-TR-84-0101

UNCLASSIFIED

F49620-80-C-0068

F/G 20/5


NL



END

DATE
FILMED

**4* 44
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AFCSE-TR- 84-0101

(3)

(to be presented at CLEO '83,
Baltimore, Maryland, May 1983).

HEPL 923

FEBRUARY 1983

AD A138683

PROGRESS AND PROBLEMS IN STORAGE RING FREE ELECTRON LASERS*

C. Bazin, M. Bergher, M. Billardon, D.A.G. Deacon, P. Elleaume, J. M. J.
Madey, J. M. Ortega, Y. Petroff, K. E. Robinson, M. P. Velghe

LURE, Universite de Paris-Sud, 91405 Orsay
France

and

High Energy Physics Laboratory
Stanford University
Stanford, California 94305

S MAR 8 1984
A

DTIC FILE COPY

Approved for public release;
distribution unlimited.

PROGRESS AND PROBLEMS IN STORAGE RING FREE ELECTRON LASERS

C. BAZIN, M. BERGHER, M. BILLARDON, D.A.G. DEACON, P. ELLEAUME, J.M.J. MADEY,
JM ORTEGA, Y. PETROFF, K.E. ROBINSON, M.F. VELGHE.

LURE, Université de Paris Sud, 91405 ORSAY, France.

and

HEPL, Stanford University, Stanford CA 94305, U.S.A.

Abstract : We discuss current problems in storage ring laser development :
optics degradation, and the low gain available on unoptimized
existing electron sources. We introduce the field with experimental
data, and conclude with the most recent results.

To be presented at CLEO '83
Baltimore, May 1983.

PROGRESS AND PROBLEMS IN STORAGE RING FREE ELECTRON LASERS

C. BAZIN, M. BERGHER, M. BILLARDON, D.A.G. DEACON, P. ELLEAUME, J.M.J. MADEY,
JM ORTEGA, Y. PETROFF, K.E. ROBINSON, M.F. VELGHE.

LURE, Université de Paris Sud, 91405 ORSAY, France.

and

HEPL, Stanford University, Stanford CA 94305, U.S.A.

A storage ring free electron laser uses an electron storage ring as the relativistic electron beam source, and an undulator magnet structure to couple the kinetic energy of the electrons into coherent laser light. Work has been underway for three years on the storage ring ACO at Orsay, France to realize a free electron laser (FEL) on a recirculated beam, with the goal of increasing the power and efficiency of the FEL. Considerable progress has been made towards the initial goal of achieving oscillation and only one problem --mirror degradation-- now stands in the way of success.

Two undulators have been studied at Orsay, the first of which had a serious magnetic field distortion problem. This problem has made possible the experimental (and theoretical) identification of the effects of electron trajectory errors on the performance of FEL's [1]. The second Orsay undulator was designed so as to eliminate these problems, and has shown detailed equivalence with the theory in its operation [2].

A modification of the undulator called the optical klystron (first operated at Novosilinsk [3]) has recently been constructed and tested at Orsay [2]. We have measured the gain with both undulators [1, 2], and more recently with the optical klystron modification. The results provide experimental confirmation of the Madey theorem [4]. The maximum peak gain per round trip in the optical cavity is 7.2×10^{-4} with the optical klystron in the Orsay system. This is an increase of a factor of five over the maximum measured undulator gain, as predicted by the theory.

AIR FORCE

NOTICE

This

approval

Distribution

MATTHEW

Chief, Technical Information Division

100-100000

has been reviewed

and approved IAW AFR 190-100

unlimited.

PER

Chief, Technical Information Division

We have obtained and measured several sets of multilayer SiO_2 and TiO_2 mirrors with total losses per round trip of less than 3×10^{-4} in air. These mirrors are inserted directly into the 10^{-10} Torr vacuum of the storage ring to eliminate window losses. Although the initial losses of the mirror pairs is sufficiently low to permit oscillation, their reflectivity degrades [2] both during insertion into the vacuum and as a result of exposure to the high flux of hard UV radiation emitted as harmonics of the undulator frequency.

We plan to test a few other mirror materials and configurations at Orsay, but a systematic attack on the mirror degradation problem is clearly needed.

The gain available on ACO is limited by the current density and straight section length. With the optical klystron in place, maximum use is made of the straight section, and the gain may be increased only by increasing the current density which depends on the magnet lattice and energy. On the other hand a new machine with high current density and a long straight section for the undulator could be expected to operate with a gain on the order of unity.

We also have measured the bunch lengthening effect [5] of the laser interaction on the stored electron beam. At low current, the magnitude and spectrum of the effect is in agreement with the theory which demonstrates that we understand at least one of the fundamental phenomena which will be responsible for saturation once the storage ring laser operates. At high the current, the coupled interactions of the "laser-induced" and the "anomalous" bunch lengthening which we have now observed indicate the need for more theoretical work.

REFERENCES

1. C. BAZIN, M. BILLARDON, D.A.G. DEACON, P. ELLEAUME, Y. FARGE, J.M.J. MADEY, J.M. ORTEGA, Y. PETROFF, K.E. ROBINSON, M.F. VELGHE, "Results of the first phase of the ACO storage ring laser experiment" in Physics of Quantum Electronics Vols 8-9, eds. S.F. JACOBS, G.T. MOORE, H.S. PILLOFF, M. SARGENT III, M.O. SCULLY, R. SPITZER, (Addison - Wesley, 1982).
2. M. BILLARDON, D.A.G. DEACON, P. ELLEAUME, J.M. ORTEGA, K.E. ROBINSON C. BAZIN, M. BERGHER, J.M.J. MADEY, Y. PETROFF, M. VELGHE, "Recent results of the ACO Storage ring FEL experiment", to be published in the Proceedings of the Bendor Free Electron Laser Conference, eds. M. BILLARDON and D.A.G. DEACON (Journal de Physique Colloques, 1983).
3. G.A. KORNYUKHIN, G.N. KULIPANOV, U.N. LITVINENKO, N.A. MEZENTSEV, A.N. SKRINSKY, N.A. VINORUROV and P.D. VOBLIY, "Experiments with an optical klystron on the VEPP-3 storage ring" (unpublished), Institute of Nuclear Physics, 630090 NOVOSIBIRSK, URSS.
4. D.A.G. DEACON, K.E. ROBINSON, J.M.J. MADEY, C. BAZIN, M. BILLARDON, P. ELLEAUME, Y. FARGE, J.M. ORTEGA, Y. PETROFF, M.F. VELGHE, Optics Communications **40**, 373 (1982).
5. K.E. ROBINSON, D.A.G. DEACON, M.F. VELGHE, J.M.J. MADEY, to be published in the IEEE Journal of Quantum Electronics (1983).

Classification For
GROUP 1
TAR
Excluded
Excluded

Classification/
Availability Codes
A-1

11-11-81

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER AFCCR-TR-84-0101	2. GOVT ACCESSION NO. AD-A138683	3. RECIPIENT'S CATALOG NUMBER	
4. TITLE (and Subtitle) PROGRESS AND PROBLEMS IN STORAGE RING FREE ELECTRON LASERS		5. TYPE OF REPORT & PERIOD COVERED INTERIM	
		6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s) C. BAZIN, M. BERGHER, M. BILLARDON, D.A.G. DEACON, P. ELLEAUME, J.M.J. MADEY, J.M. ORTEGA, Y. PETROFF, K.E. ROBINSON, M.P. VELGHE		8. CONTRACT OR GRANT NUMBER(s) F49620-80-C-0068	
9. PERFORMING ORGANIZATION NAME AND ADDRESS STANFORD UNIVERSITY, HIGH ENERGY PHYSICS LAB STANFORD, CA 94305		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2301/A1	
11. CONTROLLING OFFICE NAME AND ADDRESS AFOSR,/NP Bolling AFB, DC 20332		12. REPORT DATE FEB 83	
		13. NUMBER OF PAGES 5	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED	
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.			
17. DISTRIBUTION STATEMENT (of the abstract entered in block 20, if different from Report)			
18. SUPPLEMENTARY NOTES To be presented at CLEO '83, Baltimore, MD, May 83.			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) <i>This document discusses</i>			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We discuss present problems in storage ring laser development: optics degradation, and the laser available on unoptimized existing electron sources. We introduce the laser with experimental data, and conclude with the most recent results.			

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

END

DATE
FILMED

4-84

DTIC